

OptiPrep™ Mini-Review MS15

Endocytosis analysis – a review of density gradient methods

1. Nycodenz® gradients

1a. Long-spin Nycodenz® continuous gradients

Some of the first reports on the use of Nycodenz® for the fractionation of endosomes were published in the mid-nineteen eighties by Howard Evans and his co-workers, working at the National Institute for Medical Research in London. They worked primarily on the endocytosis of a variety of ^{125}I -labelled asialo-glycoproteins by the perfused rat liver. After differential centrifugation of the homogenate, a supernatant from a light mitochondrial pellet was first applied to a continuous 15-43% (w/v) sucrose gradient (over layers of 43% and 70% sucrose) and centrifuged at 140,000 g for 3.5 h. The lightest fraction (1.095-1.117 g/ml) was re-centrifuged on a 13.8-27.6% (w/v) Nycodenz® gradient at 110,000 g for 18 h. Two very distinct, well-separated peaks of radioactivity were obtained at 1.090 and 1.115 g/ml. The denser material was identified as early endosomes and the lighter as late endosomes [1].

1b. Short-spin Nycodenz® velocity gradients

At about the same time, Trond Berg's group at the University of Oslo, working primarily with isolated hepatocytes, developed a variety of Nycodenz® gradients for the analysis of ligand internalization. The gradients were continuous 0-45% (w/v) Nycodenz®, centrifuged at 85,000 g for a variety of times, with the sample either top- or bottom-loaded. Two principal times were chosen, either 5 h for buoyant density analysis or 45 min for rate-zonal (sedimentation or flotation velocity) analysis. The top-loaded 45 min gradients showed that, with increasing internalization time (30 sec to 30 min), asialofetuin was associated with membrane compartments of increasing sedimentation velocity [2], which were all distinct from the bulk of the lysosomes (see Figure 1). If the centrifugation was carried out for 3 h, so that the membrane vesicles reached their banding density, then there was only a relatively small difference between the peak density at 1 and 15 min incubation times (1.10 and 1.11 g/ml respectively). Using the 45 min gradient format, it was later shown, using a polyethylene glycol-modified asialofetuin, that the peak around 1.05-1.06 g/ml was early endosomes and that at 1.09 g/ml comprised multi-vesicular bodies; the latter were quite distinct from lysosomes at 1.14 g/ml [3].




Figure 1: Distribution of ^{125}I -labelled ligand at different incubation times. Also shown is the banding position of lysosomes (GNase = β -N-Acetylglucosaminidase). Figure adapted from ref 2.

1c. Hybrid Nycodenz®/polysucrose gradients

Branch et al [4] compared the efficacy of short-spin (1 h) continuous polysucrose and Nycodenz® gradients in the analysis of the membrane compartments in rat liver during the transcytosis of polymeric IgA and endocytosis of asialofetuin. In both instances a Beckman VTi50, vertical rotor was used at 206,000 g. The authors concluded that while polysucrose gradients were superior for resolving light (early) and dense (late) endosomes, Nycodenz® gradients provided far greater discrimination between late endosomes and lysosomes; moreover discrimination was achieved between lysosomes and very dense endosomes [5]. This led to the use of hybrid polysucrose-Nycodenz® gradients for the simultaneous isolation of early and late endosomes and lysosomes [6-9]; also, a simplified discontinuous gradient for the separation of lysosomes, very dense endosomes and other less dense endosomes was developed.

2. OptiPrep™

2a. Flotation velocity gradients

In 1997 the first paper reporting the use of an iodixanol gradient in endocytic studies was published by Orlandi [10]; the bottom-loaded 9-30% (w/v) iodixanol gradient centrifuged at 52,000 g for 90 min was similar to the Nycodenz® flotation velocity gradient (see Section 1b) of Kindberg et al [2]. It was used to analyze the internalization of cholera toxin by Caco-2 cells and over a period of 1 h the toxin peak showed a very clear shift from approx. 1.11 (0 min) to 1.12 (30 min) to 1.13 (60 min) g/ml.

2b. Overnight buoyant density gradients

A gradient system that has been widely used is that of Sheff et al [11]. In a study of the internalization of transferrin by MDCK cells transfected with the human Tfn receptor a shallow 5-20% (w/v) iodixanol gradient centrifuged at 100,000 g for 18 h was used to analyze a post-nuclear supernatant. Although many workers take advantage of the availability of high-performance rotors and the much lower viscosity of iodixanol gradients (compared to those of sucrose) to carry out buoyant density separations in 2-3 h, there is a widespread view that for the highest resolution of particles relatively low g-forces for long times are to be preferred. Sheff et al [11] were able to identify clearly a separation of early (EE) from recycling endosomes (RE), which were both well separated from the plasma membrane (PM) and lysosomes, which banded at lower densities (Figure 2).

It is important to note however that there may be significantly different patterns of banding in the case of other studies with other cell types. For example, in both HeLa [12] and COS-7 cells [13] EE banded at a lower density than RE, while the HeLa cell PM [12] banded close to the bottom of the gradient. LAMP-2 positive particles (late endosomes (LE) and lysosomes) from PC12 cells [14] and HeLa cells [15] banded at a higher density than the Rab5 positive EE. The banding patterns may also be influenced by practical variations in the handling of the cell homogenization.

In a study on the effect of moesin on receptor recycling in HeLa cells [16], this gradient system revealed two well separated bands of Rab7, the lighter one of which overlapped the lower density Rab5-containing vesicles (Figure 3). Barroso-González et al [16] compared control “scrambled” cells (nucleofected with scrambled oligonucleotides) and test cells nucleofected with siRNA moesin oligonucleotides. The TfR, which was located predominantly between the two Rab7 bands from control cells, shifted significantly into the low-density region in the siRNA moesin oligonucleotide-treated cells (Figure 3).

A very detailed analysis of the membranes from monkey kidney cells on an 8-25% (w/v) iodixanol gradient [17] identified (in increasing banding density) PM, lysosomes, endosomes + Golgi and two distinct bands of endoplasmic reticulum (ER). McKenzie et al [17] used the gradient to study the internalization of Shiga toxin (and its B subunit) and detected a single toxin-rich band in the Golgi + endosomes region; AlF_4^- treatment caused a pronounced dichotomy of the band. In an unrelated study Woods et al [18] also reported the resolution of two distinct fractions of ER from 3T3 cells, the denser of which was identified as perinuclear. Other uses of the long-spin continuous gradient are given in Table 1.

Table 1: Long-spin continuous gradient use

Cell type	Compartments analysed	Ref #
Neuroblastoma	EE (Rab5A), LE (Rab7); dopamine transporter	19
HepG2	EE subfractions (denser overlapping LE); transferrin labelling in denser EE	20
HeLa	Sorting/recycling endosomes, EE, LE, lysosomes; EE/Golgi cycling	21
HeLa	PM, LE; distribution of stomatin-like protein 1 and cholesterol transfer to LE	22

Idkowiak-Baldys et al [23] used a similar long-spin, low g-force strategy but started with a discontinuous 5%, 10%, 15% and 20% iodixanol gradient; although this will have become continuous during the centrifugation there may be small important differences in the final density profile when compared with a pre-formed continuous gradient. In HEK cells the PM and ER were concentrated at the light and dense ends respectively, while LAMP1-1 positive late endosomes banded about 2/3rds down the gradient and three distinct Rab11 areas were observed (Figure 4). In phorbol-ester treated cells, protein kinase C, which was broadly distributed in control cells, became markedly and distinctively concentrated in the lightest of the Rab11 bands.




Figure 2 Distribution of transferrin in MDCK cell membrane fractions in long-spin iodixanol gradient: effect of chase time (from ref 11, with kind permission of the authors and The Rockefeller University Press).




Figure 3 Distribution of markers in membranes from HeLa cells (long-spin iodixanol gradient): comparison of control and siRNA-moesin treated cells (adapted from ref 16)

- ◆ In contrast to iodixanol gradients, Percoll® gradients cannot be centrifuged for long time periods at the g -forces required for effective membrane fractionation without most of colloidal silica pelleting; thus its use in endocytosis studies is very restrictive. In studies on the trafficking of LDL cholesterol Percoll® gradients were unable to resolve PM and EE from CHO cells [24]; long-spin iodixanol gradients were able to exhibit distinctive banding patterns not only for PM and EE, but also LE and Golgi.




Figure 4: Distribution of HEK cells membrane compartments and Rab11 in iodixanol gradient: adapted from ref 23

2c. Short-spin continuous density gradients

Gradients covering more or less the same density as those described in 2b, when centrifuged for only 3 h at 100-130,000 g can provide excellent separation of EE from PM and Golgi from HeLa and HEK cells [25,26]. In the case of human HT1080 (human sarcoma) cells two LE fractions were identified; moreover chloroquine treatment of the cells induced a pronounced shift of Rab9 to the denser LE [27].

2d. Double-gradient strategy

An initial sedimentation velocity gradient followed by a second buoyant density gradient enabled Lin et al [28] to achieve very fine fractionation of light endosomes fractions from PC12 cells, in their studies on the internalization of nerve growth factor. To obtain a membrane vesicle fraction the cells were initially permeabilized by a single passage through a ball-bearing homogenizer. Semi-intact cells and larger membrane particles were removed by low-speed centrifugation. The supernatant was first layered on top of a 0-30% (w/v) iodixanol gradient and centrifuged at 133,000 g for 1.5 h. Fractions from this gradient were adjusted to 32.5% iodixanol by mixing with OptiPrep™; layered under a second 0-30% iodixanol gradient and centrifuged at the same speed for 18 h. The second fraction from the velocity gradient was resolved into two completely distinct non-overlapping sub-fractions; the lighter contained the neurotrophin receptor pTrkA plus its associated proteins APPL1 and GIPC1; the denser only contained APPL1. More recently [29] the same double-gradient strategy was able to resolve three distinct subclasses of endosomes: (1) those bearing activated receptor tyrosine kinases, (2) those containing p75NTR (tumor necrosis receptor superfamily) and (3) those exhibiting PAC1 (a G-protein-coupled receptor).




Figure 5 Distribution of labelled ligand (1 min pulse) from rat liver, at various chase times, in a self-generated iodixanol gradient.

(20 min) the ligand appears in a lysosomal or pre-lysosomal compartment.

2e. Self-generated gradients

The use of self-generated gradients was first reported by Billington et al [30] for analysis of the internalization of asialoglycoproteins by the perfused rat liver. One of its attractions is its simplicity and reproducibility of gradient formation. After homogenization of the liver, a 3000 g supernatant was simply adjusted to 12.5% (w/v) iodixanol, underlaid with 1 ml of 20% iodixanol and centrifuged in a vertical or near-vertical rotor at 350,000 g_{av} for 1.5 h.

The results of a 1 min pulse of labelled ligand after various chase times are shown in Figure 5. The data is interpreted as follows: (1 min chase) the ligand is found initially in a dense clathrin-coated vesicle; (2 min) uncoating of the vesicle reduces its density; (10 min) the ligand has been transferred to a low-density endosome and

Sometimes the gradient is generated not from an iodixanol solution of uniform density but from a discontinuous iodixanol gradient in of 10%, 20% and 30% (w/v) iodixanol (with the sample in the latter). Centrifugation for 3 h will produce a more or less linear gradient (that produced from a single concentration will tend to be shallower towards the top and steeper towards the bottom). This tri-layered gradient has an additional feature; all of the cytosolic proteins remain at the bottom of the gradient. It was used to investigate the distribution of Ra1A (a GTPase which interacts with the exocyst complex) that resides in recycling endosomes. The Ra1A in COS cells overlapped only the denser parts of the TfR and Rab11 region, but not with EEA1, syntaxin 6 or the cytosolic Akt. Indeed the best coincidence was observed with another recycling endosome marker – Rab4, which was clearly associated with a denser particle than Rab11 [31]. This gradient format has

also been used for studying the changes induced by the adenovirus early region 4 open reading frame 4 protein in 293T cells [32].

- ◆ See **Mini-Review MS-16** for a complete bibliography of all the published papers that have used gradients prepared from OptiPrep™ for the analysis of all aspects of endocytosis and trafficking within the endosomal system.
- ◆ There are several relevant OptiPrep™ Application Sheets that provide detailed protocols:
 - ◆ **Cultured cells – buoyant density gradients: Application Sheet S44**
 - ◆ **Cultured cells – self-generated gradients: Application Sheet S43**
 - ◆ **Rat liver – lysosome/late-endosome events: Application Sheet S52**
 - ◆ **Rat liver – sedimentation velocity gradients: Application Sheet S42**
 - ◆ **Rat liver – self-generated gradients: Application Sheet S43**
- ◆ All of the Application Sheets and Mini-Reviews may be accessed via the following website: www.axis-shield-density-gradient-media.com or from the flash-drive
- ◆ It is not known if all or any of the Nycodenz®-based methods can be transposed directly to iodixanol. Nycodenz® and iodixanol solutions of the same % (w/v) have approximately the same density but the former have a higher osmolality (see Application Sheets for more information).

3. References

1. Evans, W.H. and Flint, N. (1985) *Subfractions of hepatic endosomes in Nycodenz gradients and by free-flow electrophoresis* Biochem. J., **232**, 25-32
2. Kindberg, G.M., Ford, T., Blomhoff, R., Rickwood, D. and Berg, T. (1984) *Separation of endocytic vesicles in Nycodenz gradients* Anal. Biochem., **142**, 455-462
3. Roseng, I., Tolleshaug, H. and Berg, T. (1992) *Uptake, intracellular transport, and degradation of polyethylene glycol-modified asialofetuin in hepatocytes* J. Biol. Chem., **267**, 22987-22993
4. Branch, W.J., Mullock, B.M. and Luzio, J.P. (1987) *Rapid subcellular fractionation of the rat liver endocytic compartments involved in transcytosis of polymeric immunoglobulin A and endocytosis of asialofetuin* Biochem. J., **244**, 311-315
5. Perez, J. H., Branch, W. J., Smith, L., Mullock, B. M. and Luzio, J.P. (1988) *Investigation of endosomal compartments involved in endocytosis and transcytosis of polymeric immunoglobulin A by subcellular fractionation of perfused isolated rat liver* Biochem. J., **251**, 763-770
6. Mullock, B. M., Perez, J. H., Kuwana, T., Gray, S. R. and Luzio, J. P. (1994) *Lysosomes can fuse with a late endosomal compartment in a cell-free system from rat liver* J. Cell Biol., **126**, 1173-1182
7. Mullock, B. M., Bright, N. A., Fearon, C. W., Gray, S. R. and Luzio, J. P. (1998) *Fusion of lysosomes with late endosomes produces a hybrid organelle of intermediate density and is NSF dependent* J. Cell. Biol., **140**, 591-601
8. Mullock, B. M., Smith, C. W., Ihrke, G., Bright, N. A., Lindsay, M., Parkinson, E. J., Brooks, D. A. Parton, R. G., James, D. E., Luzio, J. P. and Piper, R. C. (2000) *Syntaxin 7 is localized to late endosome compartments, associates with Vamp 8, and is required for late endosome-lysosome fusion* Mol. Biol. Cell **11**, 3137-3153
9. Pryor, P. R., Mullock, B. M., Bright, N. A., Gray, S. R. and Luzio, J. P. (2000) *The role of intraorganellar Ca²⁺ in late endosome-lysosome heterotypic fusion and in the reformation of lysosomes from hybrid organelles* J. Cell Biol., **149**, 1053-1062
10. Orlandi, P.A.(1997) *Protein-disulfide isomerase-mediated reduction of the A subunit of cholera toxin in a human intestinal cell line* J. Biol. Chem., **272**, 4591-4599
11. Sheff, D.R., Daro, E.A., Hull, M. and Mellmann, I. (1999) *The receptor recycling pathway contains two distinct populations of early endosomes with different sorting functions* J. Cell Biol., **145**, 123-139
12. Meyers, J.M. and Prekeris, R. (2002) *Formation of mutually exclusive Rab11 complexes with members of the family of Rab11-interacting proteins regulates Rab11 endocytic targeting and function* J. Biol. Chem., **277**, 49003-49010
13. Shen, X., Xu, K-F., Fan, Q., Pacheco-Rodriguez, G., Mos, J. and Vaughan, M. (2006) *Association of brefeldin A-inhibited guanine nucleotide-exchange protein 2 (BIG2) with recycling endosomes during transferring uptake* Proc. Natl. Acad. Sci. USA, **103**, 2635-2640
14. Li, Y., Chin, L-S., Levey, A.L. and Li, L. (2002) *Huntingtin-associated protein 1 interacts with hepatocyte growth factor-regulated tyrosine kinase substrate and functions in endosomal trafficking* J. Biol. Chem., **277**, 28212-28221
15. Chin, L-S., Raynor, M.C., Wei, X., Chen, H-Q. and Li, L. (2001) *Hrs interacts with sorting nexin 1 and regulates degradation of epidermal growth factor receptor* J. Biol. Chem., **276**, 7069-7078

16. Barroso-González, J., Machado, J-D., García-Expósito, L. and Valenzuela-Fernández, A. (2009) *Moesin regulates the trafficking of nascent clathrin-coated vesicles* J. Biol. Chem., **284**, 2419–2434
17. McKenzie, J., Johannes, L., Taguchi, T. and Sheff, D. (2009) *Passage through the Golgi is necessary for Shiga toxin B subunit to reach the endoplasmic reticulum* FEBS J., **276**, 1581–1595
18. Woods, A.J., Roberts, M.S., Choudhary, J., Barry, S.T., Mazaki, Y., Sabe, H., Morley, S.J., Critchley, D.R. and Norman, J.C. (2002) *Paxillin associates with poly(A)-binding protein 1 at the dense endoplasmic reticulum and the leading edge of migrating cells* J. Biol. Chem., **277**, 6428–6437
19. Wiesinger, J.A., Buwen, J.P., Cifelli, C.J., Unger, E.L., Jones, B.C. and Beard, J.L. (2007) *Down-regulation of dopamine transporter by iron chelation in vitro is mediated by altered trafficking, not synthesis* J. Neurochem., **100**, 167–179
20. Manunta, M., Izzo, L., Duncan, R. and Jones, A.T. (2007) *Establishment of subcellular fractionation techniques to monitor the intracellular fate of polymer therapeutics II: Identification of endosomal and lysosomal compartments in HepG2 cells combining single-step subcellular fractionation and fluorescent imaging* J. Drug Target., **15**, 37–50
21. Proikas-Cezanne, T., Gaugel, A., Frickey, T. and Nordheim, A. (2006) *Rab14 is part of the early endosomal clathrin-coated TGN microdomain* FEBS Lett., **580**, 5241–5246
22. Mairhofer, M., Steiner, M., Salzer, U. and Prohaska, R. (2009) *Stomatin-like protein-1 interacts with stomatin and is targeted to late endosomes* J. Biol. Chem., **284**, 29218–29229
23. Idkowiak-Baldys, J., Becker, K.P., Kitatani, K. and Hannum, Y.A. (2006) *Dynamic sequestration of the recycling compartment by classical protein kinase C* J. Biol. Chem., **281**, 22321–22331
24. Sugii, S., Reid, P.C., Ohgami, N., Du, H. and Chang, T-Y. (2003) *Distinct endosomal compartments in early trafficking of low density lipoprotein-derived cholesterol* J. Biol. Chem., **278**, 27180–27189
25. Tagami, S., Okochi, M., Yanagida, K., Ikuta, A., Fukumori, A., Matsumoto, N., Ishizuka-Katsura, Y., Nakayama, T., Itoh, N., Jiang, J., Nishitomi, K., Kamino, K., Morihara, T., Hashimoto, R., Tanaka, T., Kudo, T., Chiba, S. and Takeda, M. (2008) *Regulation of Notch signaling by dynamic changes in the precision of S3 cleavage of Notch-1* Mol. Cell. Biol., **28**, 165–76
26. Fukumori, A., Okochi, M., Tagami, S., Jiang, J., Itoh, N., Nakayama, T., Yanagida, K., Ishizuka-Katsura, Y., Morihara, T., Kamino, K., Tanaka, T., Kudo, T., Tanii, H., Ikuta, A., Haass, C. and Takeda, M. (2006) *Presenilin-dependent γ -secretase on plasma membrane and endosomes is functionally distinct* Biochemistry, **45**, 4907–4914
27. Molle, D., Segura-Morales, C., Camus, G., Berlioz-Torrent, C., Kjems, J., Basyuk, E. and Bertrand, E. (2009) *Endosomal trafficking of HIV-1 Gag and genomic RNAs regulates viral egress* J. Biol. Chem., **284**, 19727–19743
28. Lin, D.C., Quevedo, C., Brewer, N.E., Bell, A., Testa, J.R., Grimes, M.L., Miller, F.D. and Kaplan, D.R. (2006) *APPL1 associates with TrkA and GIPC1 and is required for nerve growth factor-mediated signal transduction* Mol. Cell. Biol., **26**, 8928–8941
29. McCaffrey, G., Welker, J., Scott, J., van der Salm, L. and Grimes, M.L. (2009) *High-resolution fractionation of signaling endosomes containing different receptors* Traffic, **10**, 938–950
30. Billington, D., Maltby, P.J., Jackson, A.P. and Graham, J.M. (1998) *Dissection of hepatic receptor-mediated endocytic pathways using self-generated gradients of iodixanol (OptiPrep)* Anal. Biochem., **258**, 251–258
31. Chen, X-W., Inoue, M., Hsu, S. and Saltiel, A.R. (2006) *RalA-exocyst-dependent recycling endosome trafficking is required for the completion of cytokinesis* J. Biol. Chem., **281**, 38609–38616
32. Landry, M-C., Sicotte, A., Champagne, C. and Lavoie, J.N. (2009) *Regulation of cell death by recycling endosomes and Golgi membrane dynamics via a pathway involving Src-family kinases, Cdc42 and Rab11a* Mol Biol. Cell, **20**, 4091–4106

Mini-Review MS15 5th edition, January 2018

Alere Technologies AS

Axis-Shield Density Gradient Media
is a brand of Alere Technologies AS